

Massless Higher Spin Fields in the AdS Background and BRST Constructions for Nonlinear Algebras

I.L. Buchbinder^{a *}, A. Pashnev^{b†} and M. Tsulaia^{b,c‡}

^a*Department Theoretical Physics, Tomsk State Pedagogical University
Tomsk, 634041, Russia*

^b*Bogoliubov Laboratory of Theoretical Physics, JINR
Dubna, 141980, Russia*

^c*The Andronikashvili Institute of Physics, Georgian Academy of Sciences,
Tbilisi, 380077, Georgia*

Abstract

The detailed description of the method of the construction of the nilpotent BRST charges for nonlinear algebras of constraints appearing in the description of the massless higher spin fields on the AdS_D background is presented. It is shown that the corresponding BRST charge is not uniquely defined, but this ambiguity has no impact on the physical content of the theory.

BRST – BFV method [1] is a powerful tool for the quantization of the wide class of the physical systems, invariant with respect to the gauge symmetry groups. The physical content of the theory is singled out by the BRST quantization condition

$$Q|Phys\rangle = 0 \quad (1)$$

for the nilpotent $Q^2 = 0$ BRST charge. Obviously due to the nilpotence property of the BRST charge the BRST quantization condition is gauge invariant i.e., the physical states are defined up to the transformation $|Phys\rangle' = |Phys\rangle + Q|\Lambda\rangle$ where the last term is a “spurious” state with the zero norm. Therefore to find the physical spectrum of the theory, one has to solve a cohomology problem of the corresponding BRST charge.

On the other hand the BRST method can be applied for the construction of the field theory, which corresponds to some first quantized system. Namely if one succeeds to construct the nilpotent BRST charge for the given system of constraints, then one can straightforwardly write the corresponding field theoretical Lagrangian of the form $L \sim \langle \Phi|Q|\Phi \rangle$ which gives the condition (1) as the equation of motion. At the same time the Lagrangian is gauge invariant under the transformations

$$\delta|\Phi\rangle = Q|\Lambda\rangle, \quad (2)$$

The later property is crucial for eliminating nonphysical degrees of freedom from the spectrum. The famous example of such kind of construction is BRST string field theory.

The method of construction of the nilpotent BRST charge is well known when the corresponding constraints form the closed linear Lie algebra. The situation becomes more

*E-mail: joseph@tspu.edu.ru

†E-mail: pashnev@thsun1.jinr.dubna.su

‡E-mail: tsulaia@thsun1.jinr.ru

complicated when one deals with the second – class constraints and (or) the algebra of constraints is nonlinear. In the present paper we analyze this situation and show, that at least for the system of constraints describing irreducible massless higher spin fields the corresponding BRST charge is not unique. Rather it can contain a number of free parameters which, nevertheless does not affect the physical content of the corresponding field theory. These parameters, leading to the noncanonical form of the BRST charge, can be present even in the case of closed linear algebra of constraints. The particular choice of these parameters ensures the possibility to simplify the expression for the BRST charge, and correspondingly the procedure of construction of the field theoretical Lagrangian becomes simpler as well.

Below we concentrate on the construction of the nilpotent BRST charges and corresponding field theoretical Lagrangian for the theory describing the higher spin fields in the flat space and in AdS spaces. In particular, we investigate an arbitrariness in the BRST charge, which is described by some free parameters, and show that the physical content of the corresponding field theory is independent of the values of these parameters.

First we briefly describe the system of the free higher massless spin fields propagating through the D – dimensional flat Minkowski space time in the framework of the BRST approach [2] and then generalize the method for the interaction with AdS_D background [3].

For the free massless higher spin fields the corresponding system of constraints imposed on the physical states includes the mass – shell constraint $L_0 = p_\mu^2$, transversality condition $L_1 = p_\mu a^\mu$, and tracelessness condition $L_2 = \frac{1}{2} a_\mu a^\mu$, where

$$p_\mu = \partial_\mu, \quad [a_\alpha, a_\beta^+] = \eta_{\alpha\beta}, \quad \eta_{\alpha\beta} = \text{diag}(-1, 1, 1, \dots, 1), \quad (3)$$

These constraints are written down in terms of creation and annihilation operators a_μ^+, a^μ . The general vector in the Fock space generated by these operators has the form

$$|\Phi\rangle = \sum_n \Phi_{\alpha_1 \alpha_2 \dots \alpha_n}(x) a^{\alpha_1+} a^{\alpha_2+} \dots a^{\alpha_n+} |0\rangle. \quad (4)$$

Obviously, the high spin fields $\Phi_{\alpha_1 \alpha_2 \dots \alpha_n}(x)$ are automatically symmetrical with respect to the permutation of their indices and therefore representations of the corresponding flat space little group $O(d-2)$ are characterized by the Young tableaux with one row. Note, that more complicated Young tableaux need for their description more then one set of creation and annihilation operators.

The operators L_0, L_1, L_2 along with their conjugates L_1^+ and L_2^+ obey the following commutation relations.

$$\begin{aligned} [L_1^+, L_1] &= -L_0, & [L_1^+, L_2] &= -L_1, & [L_2^+, L_1] &= -L_1^+, \\ [G_0, L_1] &= -L_1, & [L_1^+, G_0] &= -L_1^+, \end{aligned} \quad (5)$$

and

$$[G_0, L_2] = -2L_2, \quad [L_2^+, G_0] = -2L_2^+, \quad [L_2^+, L_2] = -G_0. \quad (6)$$

where we have denoted $G_0 \equiv a^{+\mu} a_\mu + \frac{D}{2}$.

The construction of the nilpotent BRST charge for this system of constraints is not straightforward since as it can be seen from (6) the constraints $L_2^\pm$ are of the second class. It is a consequence of the fact that the operator G_0 is strictly positive and therefore can not be considered as an additional constraint.

Let us first introduce the anticommuting ghost variables $\eta_0, \eta_1^+, \eta_1, \eta_2^+, \eta_2, \eta_G$ which correspond to the operators $L_0, L_1, L_1^+, L_2, L_2^+$ and G_0 and have the ghost number equal to 1, as well as the corresponding momenta $\mathcal{P}_0, \mathcal{P}_1, \mathcal{P}_1^+, \mathcal{P}_2, \mathcal{P}_2^+, \mathcal{P}_G$ with the ghost number -1 obeying the anticommutation relations

$$\{\eta_0, \mathcal{P}_0\} = \{\eta_2, \mathcal{P}_2^+\} = \{\eta_2^+, \mathcal{P}_2\} = \{\eta_2, \mathcal{P}_2^+\} = \{\eta_2^+, \mathcal{P}_2\} = \{\eta_G, \mathcal{P}_G\} = 1. \quad (7)$$

Then we build the auxiliary representations of the generators of the algebra (6) in terms of the additional creation and annihilation operators b^+ and b , $[b, b^+] = 1$ with the help of the following construction using the Verma module. We introduce the vector in the space of the Verma module $|n\rangle_V = (L_2^+)^n |0\rangle_V$, $n \in \mathbb{N}$, $L_2 |0\rangle_V = 0$ and the corresponding vector in the Fock space

$$|n\rangle = (b^+)^n |0\rangle, \quad b|0\rangle = 0.$$

Mapping the vector in the space of Verma module into the vector in the Fock space one obtains the representations of the generators of algebra $SO(2, 1)$ in terms of the variables b^+ and b

$$L_{2.aux}^+ = b^+, \quad G_{0.aux} = 2b^+b + h, \quad L_{2.aux} = b^+bb + bh \quad (8)$$

where h is the highest weight of the Verma module $G_{0.aux}|0\rangle = h|0\rangle$. Then defining the operators $\tilde{L}_2^\pm = L_2^\pm + L_{2.aux}^\pm$, $\tilde{G}_0 = G_0 + G_{0.aux}$, we construct the corresponding “bare” nilpotent BRST charge for the total system of constraints (5)–(6)

$$\begin{aligned} \tilde{Q} = & \eta_0 L_0 + \eta_1 L_1^+ + \eta_1^+ L_1 + \eta_2 \tilde{L}_2^+ + \eta_2^+ \tilde{L}_2 + \\ & + \eta_G (-\tilde{G}_0 + 3 - \eta_1^+ \mathcal{P}_1 - \mathcal{P}_1^+ \eta_1 - 2\mathcal{P}_2^+ \eta_2 - 2\eta_2^+ \mathcal{P}_2) \\ & + \eta_2^+ \eta_2 \mathcal{P}_G - \eta_1^+ \eta_1 \mathcal{P} + \eta_1^+ \mathcal{P}_1^+ \eta_2 - \eta_2^+ \eta_1 \mathcal{P}_1. \end{aligned} \quad (9)$$

In order to avoid the condition of the type $G_0|\Phi\rangle = 0$ one has to get rid of the variables η_G and $\mathcal{P}_G$ in the BRST charge (9), keeping its nilpotence property at the same time. This can be done by replacing the parameter h by the expression

$$\pi = -(G_0 + 2b^+b - 3 + \eta_1^+ \mathcal{P}_1 + \mathcal{P}_1^+ \eta_1 + 2\mathcal{P}_2^+ \eta_2 + 2\eta_2^+ \mathcal{P}_2) \quad (10)$$

and then simply omit the dependence on $\mathcal{P}_G$ in the BRST charge (see [4] – [5] for the general treatment).

The “reduced” BRST charge thus takes the form

$$\begin{aligned} Q = & \eta_0 L_0 + \eta_2 (L_2^+ + b^+) + \eta_1 L_1^+ + \eta_1^+ L_1 + \eta_2^+ (L_2 - G_0 b + b - b^+ b b) \\ & - \eta_1^+ \eta_1 \mathcal{P} + \eta_1^+ \mathcal{P}_1^+ \eta_2 - \eta_2^+ \eta_1 \mathcal{P}_1 + \eta_1^+ \eta_2^+ \mathcal{P}_1 b - \eta_2^+ \mathcal{P}_1^+ \eta_1 b - 2\eta_2^+ \mathcal{P}_2^+ \eta_2 b \end{aligned} \quad (11)$$

Finally to restore the hermiticity property of the BRST charge, which is lost due to the nonhermitian form of auxiliary representations (8) one has to define the scalar product in the Fock space as $\langle \Phi_1 | K | \Phi_2 \rangle$ where K is a nondegenerate kernel operator

$$K = Z^+ Z, \quad Z = \sum_n \frac{1}{n!} (L_2^+)^n |0\rangle_V \langle 0| (b)^n \quad (12)$$

satisfying the property $KQ = Q^+ K$.

Let us define the ghost vacuum in the way that the operators $\eta_0, \eta_1^+, \eta_2^+, \mathcal{P}_1^+$ and $\mathcal{P}_2^+$ are creation operators. Then the BRST invariant Lagrangian for our system will be.

$$L = \int \eta_0 \langle \chi | K Q | \chi \rangle \quad (13)$$

The state vector $|\chi\rangle$ along with the oscillators a_μ^+ and b^+ depends on the ghost creation operators as well and therefore the Lagrangian (13) contains some auxiliary fields. However one can show that after making use of the BRST gauge invariance (2) and the equations of motion with respect to the auxiliary fields, one is left with the only physical field with no ghost and b^+ dependence. This field is double traceless $L_2 L_2 |\Phi\rangle = 0$ and the corresponding Lagrangian for this field has the form [6]

$$L = \langle \Phi | L_0 - L_1^+ L_1 - 2L_2^+ L_0 L_2 + L_2^+ L_1 L_1 + L_1^+ L_1^+ L_2 - L_2^+ L_1^+ L_1 L_2 | \Phi \rangle \quad (14)$$

and the Lagrangian (14), is invariant under the gauge transformations

$$\delta |\Phi\rangle = L_1^+ |\lambda_1\rangle \quad (15)$$

with the traceless parameter of gauge transformations $L_2 |\lambda_1\rangle = 0$.

Let us note that, the “bare” nilpotent BRST charge (9) has been constructed in the standard fashion adopted for the system of the first – class constraints i.e., the terms nonlinear in ghost variables have the form given in [1]. However there exists an ambiguity in the expression of the BRST charge for this system as we shall see below.

Now we generalize this procedure for the description of the higher massless integer spin fields interacting with D – dimensional Anti - de - Sitter background. The main feature we face in this case is the nonlinear character of the corresponding algebra of constraints [7]. In the AdS space the momentum operator p_μ (here we denote Einstein indexes as $\mu, \nu, \dots$ and the tangent space indexes as $\alpha, \beta, \dots$) gets modified as

$$p_\mu = \partial_\mu + \omega_\mu^{\alpha\beta} a_\alpha^+ a_\beta. \quad (16)$$

Using this equation one can easily check that the operators $L_1^\pm = a^{\mu\pm} p_\mu$ are mutually hermitian conjugated with respect to the integration measure $d^D x \sqrt{-g}$. After introducing the covariant d’Alambertian

$$L_0 = g^{\mu\nu} (p_\mu p_\nu - \Gamma_{\mu\nu}^\lambda p_\lambda) \quad (17)$$

one obtains along with (6) the following commutation relations

$$\begin{aligned} [L_1^+, L_1] &= -\tilde{L}_0, & [L_1^+, L_2] &= -L_1, & [L_2^+, L_1] &= -L_1^+, \\ [\tilde{L}_0, L_1] &= -2rL_1 + 4rG_0L_1 - 8rL_1^+L_2, \\ [L_1^+, \tilde{L}_0] &= -2rL_1^+ + 4rL_1^+G_0 - 8rL_2^+L_1, \\ [G_0, L_1] &= -L_1, & [L_1^+, G_0] &= -L_1^+, \end{aligned} \quad (18)$$

where

$$\tilde{L}_0 \equiv L_0 + r(-D + \frac{D^2}{4}) + 4rL_2^+L_2 - rG_0G_0 + 2rG_0, \quad (19)$$

and the parameter r being related to the inverse radius of the AdS_D space λ via $r = \lambda^2$.

As it can be seen from the relations (18) and (6) the operators obey the nonlinear algebra being analogous to the finite $W_3^{(2)}$ algebra [8]. Again our goal is to construct the nilpotent BRST charge for this system, where the operator G_0 is excluded from the total set of constraints.

As a first step we construct nilpotent BRST charge for the operators $L_0, L_1^\pm, L_2^\pm$ and G_0 satisfying the nonlinear algebra (6), (18). However in contrast to the case of the

flat space – time i.e., when the constraints form the linear algebra there is no standard prescription for the construction of nilpotent BRST charge for the system of operators forming a nonlinear algebra. For this end we add step by step all possible higher order terms in the ghosts with arbitrary coefficients to the BRST charge (9) and require its nilpotence. Simultaneously we modify the operator at the linear term in ghost variable η_0 , instead of being simply L_0 it becomes the nonlinear combination $\tilde{L}_0$ of L_0 , G_0 and $L_2^\pm$ (19).

This procedure leads to the family of “bare” nilpotent BRST charges, which turn out to depend on three free parameters k_1, k_2, k_3 , namely i.e, the expression of terms which contain higher degrees in the ghost variables turn out not to be uniquely defined. The possible solution of the problem (not necessarily the most general one) has the form

$$\tilde{Q}^1 = \tilde{Q}_0^1 + k_1 \tilde{Q}_{k_1}^1 + k_2 \tilde{Q}_{k_2}^1 + k_3 \tilde{Q}_{k_3}^1, \quad (20)$$

where

$$\begin{aligned} \tilde{Q}_0^1 = & \eta_0(\tilde{L}_0 + 6r) + \eta_1 L_1^+ + \eta_2 L_2^+ + \eta_1^+ L_1 + \eta_2^+ L_2 - \eta_G(G_0 - 3) \\ & + 2r\eta_0\eta_1^+ \mathcal{P}_1 - 8r\eta_0\eta_2^+ \mathcal{P}_2 + 2r\eta_0\mathcal{P}_1^+ \eta_1 \\ & - 8r\eta_0\mathcal{P}_2^+ \eta_2 - \eta_1^+ \eta_1 \mathcal{P}_0 + \eta_1^+ \eta_G \mathcal{P}_1 + \eta_1^+ \mathcal{P}_1^+ \eta_2 - 4r\eta_1^+ \mathcal{P}_G \eta_1 \\ & - \eta_2^+ \eta_1 \mathcal{P}_1 + 2\eta_2^+ \eta_G \mathcal{P}_2 - \eta_2^+ \mathcal{P}_G \eta_2 + \mathcal{P}_1^+ \eta_G \eta_1 + 2\mathcal{P}_2^+ \eta_G \eta_2 - 8r\eta_0\eta_1 \mathcal{P}_1 L_2^+ \\ & - 4r\eta_0\eta_1^+ \mathcal{P}_1 G_0 + 8r\eta_0\eta_1^+ \mathcal{P}_1^+ L_2 - 4r\eta_0\mathcal{P}_1^+ \eta_1 G_0 - 12r\eta_0\eta_1^+ \mathcal{P}_1^+ \eta_1 \mathcal{P}_1 \end{aligned} \quad (21)$$

$$\begin{aligned} \tilde{Q}_{k_1}^1 = & -2\eta_0(G_0 - 3) - 6\eta_0\eta_2^+ \mathcal{P}_2 - 6\eta_0\mathcal{P}_2^+ \eta_2 - 3\eta_1^+ \mathcal{P}_G \eta_1 - 2\eta_0\eta_1 \mathcal{P}_1 L_2^+ \\ & - \eta_0\eta_1^+ \mathcal{P}_1 G_0 - 2\eta_0\eta_1^+ \mathcal{P}_2 L_1^+ + 2\eta_0\eta_1^+ \mathcal{P}_1^+ L_2 - \eta_0\eta_1^+ \mathcal{P}_G L_1 \\ & - \eta_0\mathcal{P}_1^+ \eta_1 G_0 - 2\eta_0\mathcal{P}_2^+ \eta_1 L_1 - \eta_0\mathcal{P}_G \eta_1 L_1^+ - 6\eta_0\eta_1^+ \mathcal{P}_1^+ \eta_1 \mathcal{P}_1, \end{aligned} \quad (22)$$

$$\tilde{Q}_{k_2}^1 = -\eta_0(G_0 - 3) - \eta_0\eta_1^+ \mathcal{P}_1 - 2\eta_0\eta_2^+ \mathcal{P}_2 - \eta_0\mathcal{P}_1^+ \eta_1 - 2\eta_0\mathcal{P}_2^+ \eta_2 - \eta_1^+ \mathcal{P}_G \eta_1 \quad (23)$$

$$\begin{aligned} \tilde{Q}_{k_3}^1 = & \eta_0\mathcal{P}_2^+ \eta_2 + \eta_0\eta_2^+ \mathcal{P}_2 + \eta_0\eta_1 \mathcal{P}_1 L_2^+ + \eta_0\eta_1^+ \mathcal{P}_2 L_1^+ - \eta_0\eta_1^+ \mathcal{P}_1^+ L_2 + \eta_0\mathcal{P}_2^+ \eta_1 L_1 \\ & + \eta_0\eta_1^+ \eta_2^+ \mathcal{P}_1 \mathcal{P}_2 + 2\eta_0\eta_1^+ \mathcal{P}_1^+ \eta_1 \mathcal{P}_1 + \eta_0\eta_1^+ \mathcal{P}_1^+ \mathcal{P}_G \eta_2 - \eta_0\eta_1^+ \mathcal{P}_2^+ \eta_2 \mathcal{P}_1 \\ & - \eta_0\eta_2^+ \mathcal{P}_1^+ \eta_1 \mathcal{P}_2 - \eta_0\eta_2^+ \mathcal{P}_G \eta_1 \mathcal{P}_1 + \eta_0\mathcal{P}_1^+ \mathcal{P}_2^+ \eta_1 \eta_2 \end{aligned} \quad (24)$$

All these operators are nilpotent and mutually anticommuting so that their sum is nilpotent as well. Let us note that the particular choice of the parameters $k_1 = 0, k_2 = 0, k_3 = 8$ leads to the BRST charge constructed in [3], which includes terms up to the fifth order in ghosts. The simplest expression including only third powers of ghosts corresponds to the choice $k_1 = -2, k_2 = 4, k_3 = 0$:

$$\begin{aligned} \tilde{Q}^1 = & \eta_0(\tilde{L}_0 + 6r) + \eta_1 L_1^+ + \eta_2 L_2^+ + \eta_1^+ L_1 + \eta_2^+ L_2 \\ & - \eta_G(G_0 - 3) - 2r\eta_0\eta_1^+ \mathcal{P}_1 - 4r\eta_0\eta_2^+ \mathcal{P}_2 \\ & - 2r\eta_0\mathcal{P}_1^+ \eta_1 - 4r\eta_0\mathcal{P}_2^+ \eta_2 - \eta_1^+ \eta_1 \mathcal{P}_0 + \eta_1^+ \eta_G \mathcal{P}_1 + \eta_1^+ \mathcal{P}_1^+ \eta_2 - 2r\eta_1^+ \mathcal{P}_G \eta_1 \\ & - \eta_2^+ \eta_1 \mathcal{P}_1 + 2\eta_2^+ \eta_G \mathcal{P}_2 - \eta_2^+ \mathcal{P}_G \eta_2 + \mathcal{P}_1^+ \eta_G \eta_1 + 2\mathcal{P}_2^+ \eta_G \eta_2 - 4r\eta_0\eta_1 \mathcal{P}_1 L_2^+ \\ & - 2r\eta_0\eta_1^+ \mathcal{P}_1 G_0 + 4r\eta_0\eta_1^+ \mathcal{P}_2 L_1^+ + 4r\eta_0\eta_1^+ \mathcal{P}_1^+ L_2 + 2r\eta_0\eta_1^+ \mathcal{P}_G L_1 \\ & - 2r\eta_0\mathcal{P}_1^+ \eta_1 G_0 + 4r\eta_0\mathcal{P}_2^+ \eta_1 L_1 + 2r\eta_0\mathcal{P}_G \eta_1 L_1^+. \end{aligned} \quad (25)$$

To get rid of the nonphysical constraint G_0 we again take the auxiliary representations for the operators $L_2^\pm$ and G_0 in the form (8) and construct the “bare” nilpotent BRST

charge adding step by step all possible higher order terms in the ghosts. The result is similar to (20), namely

$$\tilde{Q} = \tilde{Q}_0 + k_1 \tilde{Q}_{k_1} + k_2 \tilde{Q}_{k_2} + k_3 \tilde{Q}_{k_3}, \quad (26)$$

but now

$$\begin{aligned} \tilde{Q}_0 = & \eta_0(\tilde{L}_0 + 6r - 4rG_{0.aux}) + \eta_1 L_1^+ + \eta_1^+ L_1 \\ & + \eta_2(L_{2.aux}^+ + L_2^+) + \eta_2^+(L_{2.aux} + L_2) - \eta_G(G_{0.aux} + G_0 - 3) \\ & + 2r\eta_0\eta_1^+\mathcal{P}_1 - 8r\eta_0\eta_2^+\mathcal{P}_2 + 2r\eta_0\mathcal{P}_1^+\eta_1 \\ & - 8r\eta_0\mathcal{P}_2^+\eta_2 - \eta_1^+\eta_1\mathcal{P}_0 + \eta_1^+\eta_G\mathcal{P}_1 + \eta_1^+\mathcal{P}_1^+\eta_2 - 4r\eta_1^+\mathcal{P}_G\eta_1 - \eta_2^+\eta_1\mathcal{P}_1 \\ & + 2\eta_2^+\eta_G\mathcal{P}_2 - \eta_2^+\mathcal{P}_G\eta_2 + \mathcal{P}_1^+\eta_G\eta_1 + 2\mathcal{P}_2^+\eta_G\eta_2 - 8r\eta_0\eta_1\mathcal{P}_1L_2^+ \\ & - 4r\eta_0\eta_1^+\mathcal{P}_1G_0 + 8r\eta_0\eta_1^+\mathcal{P}_1^+L_2 - 4r\eta_0\mathcal{P}_1^+\eta_1G_0 - 12r\eta_0\eta_1^+\mathcal{P}_1^+\eta_1\mathcal{P}_1, \end{aligned} \quad (27)$$

$$\begin{aligned} \tilde{Q}_{k_1} = & -2\eta_0(G_{0.aux} + G_0 - 3) - 6\eta_0\eta_2^+\mathcal{P}_2 - 6\eta_0\mathcal{P}_2^+\eta_2 - 3\eta_1^+\mathcal{P}_G\eta_1 \\ & - 2\eta_0\eta_1\mathcal{P}_1(L_{2.aux}^+ + L_2^+) + 2\eta_0\eta_1^+\mathcal{P}_1^+(L_{2.aux} + L_2) - \eta_0\eta_1^+\mathcal{P}_1(G_{0.aux} + G_0) \\ & - 2\eta_0\eta_1^+\mathcal{P}_2L_1^+ - \eta_0\eta_1^+\mathcal{P}_GL_1 - \eta_0\mathcal{P}_1^+\eta_1(G_{0.aux} + G_0) \\ & - 2\eta_0\mathcal{P}_2^+\eta_1L_1 - \eta_0\mathcal{P}_G\eta_1L_1^+ - 6\eta_0\eta_1^+\mathcal{P}_1^+\eta_1\mathcal{P}_1, \end{aligned} \quad (28)$$

$$\begin{aligned} \tilde{Q}_{k_2} = & -\eta_0(G_{0.aux} + G_0 - 3) - \eta_0\eta_1^+\mathcal{P}_1 - 2\eta_0\eta_2^+\mathcal{P}_2 \\ & - \eta_0\mathcal{P}_1^+\eta_1 - 2\eta_0\mathcal{P}_2^+\eta_2 - \eta_1^+\mathcal{P}_G\eta_1, \end{aligned} \quad (29)$$

$$\begin{aligned} \tilde{Q}_{k_3} = & \eta_0\eta_2^+\mathcal{P}_2 + \eta_0\mathcal{P}_2^+\eta_2 + \eta_0\eta_1\mathcal{P}_1(L_{2.aux}^+ + L_2^+) - \eta_0\eta_1^+\mathcal{P}_1^+(L_{2.aux} + L_2) \\ & + \eta_0\eta_1^+\mathcal{P}_2L_1^+ + \eta_0\mathcal{P}_2^+\eta_1L_1 + \eta_0\eta_1^+\eta_2^+\mathcal{P}_1\mathcal{P}_2 + 2\eta_0\eta_1^+\mathcal{P}_1^+\eta_1\mathcal{P}_1 \\ & + \eta_0\eta_1^+\mathcal{P}_1^+\mathcal{P}_G\eta_2 - \eta_0\eta_1^+\mathcal{P}_2^+\eta_2\mathcal{P}_1 - \eta_0\eta_2^+\mathcal{P}_1^+\eta_1\mathcal{P}_2 \\ & - \eta_0\eta_2^+\mathcal{P}_G\eta_1\mathcal{P}_1 + \eta_0\mathcal{P}_1^+\mathcal{P}_2^+\eta_1\eta_2. \end{aligned} \quad (30)$$

Taking the free parameters as in (25) again leads to the simplest form of the BRST charge

$$\begin{aligned} \tilde{Q} = & \eta_0(\tilde{L}_0 + 6r - 4rG_{0.aux}) + \eta_1 L_1^+ + \eta_1^+ L_1 \\ & + \eta_2(L_{2.aux}^+ + L_2^+) + \eta_2^+(L_{2.aux} + L_2) - \eta_G(G_{0.aux} + G_0 - 3) \\ & - 2r\eta_0\eta_1^+\mathcal{P}_1 - 4r\eta_0\eta_2^+\mathcal{P}_2 - 2r\eta_0\mathcal{P}_1^+\eta_1 - 4r\eta_0\mathcal{P}_2^+\eta_2 - \eta_1^+\eta_1\mathcal{P}_0 \\ & + \eta_1^+\eta_G\mathcal{P}_1 + \eta_1^+\mathcal{P}_1^+\eta_2 - 2r\eta_1^+\mathcal{P}_G\eta_1 - \eta_2^+\eta_1\mathcal{P}_1 \\ & + 2\eta_2^+\eta_G\mathcal{P}_2 - \eta_2^+\mathcal{P}_G\eta_2 + \mathcal{P}_1^+\eta_G\eta_1 + 2\mathcal{P}_2^+\eta_G\eta_2 + 4r\eta_0\eta_1\mathcal{P}_1L_{2.aux}^+ \\ & - 4r\eta_0\eta_1\mathcal{P}_1L_2^+ + 2r\eta_0\eta_1^+\mathcal{P}_1G_{0.aux} - 2r\eta_0\eta_1^+\mathcal{P}_1G_0 + 4r\eta_0\eta_1^+\mathcal{P}_2L_1^+ \\ & - 4r\eta_0\eta_1^+\mathcal{P}_1^+L_{2.aux} + 4r\eta_0\eta_1^+\mathcal{P}_1^+L_2 + 2r\eta_0\eta_1^+\mathcal{P}_G\mathcal{P}_1 + 2r\eta_0\mathcal{P}_1^+\eta_1G_{0.aux} \\ & - 2r\eta_0\mathcal{P}_1^+\eta_1G_0 + 4r\eta_0\mathcal{P}_2^+\eta_1L_1 + 2r\eta_0\mathcal{P}_G\eta_1L_1^+. \end{aligned} \quad (31)$$

The further procedure goes on in the complete analogy with the case of the flat space – time background. Namely we obtain the reduced BRST charges Q from (26) after the transformation (10) and then construct the Lagrangian (13) being invariant under the gauge transformation (2). Though the BRST charge which corresponds to the system (18), (6) is not unique, one can show following the lines of [3] on the base of the explicit calculations that all these BRST charges after making the partial BRST gauge fixing in the Lagrangian (13) lead to the unique final form of the Lagrangian, which contains only one double traceless physical field $|\Phi\rangle$ ($L_2L_2|\Phi\rangle = 0$)

$$\begin{aligned} \mathcal{L} = & \langle \Phi | \tilde{L}_0 - L_1^+ L_1 - 2L_2^+ \tilde{L}_0 L_2 + L_2^+ L_1 L_1 + L_1^+ L_1^+ L_2 - L_2^+ L_1^+ L_1 L_2 \\ & - r(6 - 4G_0 + 10L_2^+ L_2 - 4L_2^+ G_0 L_2) | \Phi \rangle. \end{aligned} \quad (32)$$

The Lagrangian is invariant with respect to the gauge transformation (15) with the traceless parameter $|\lambda\rangle$ and describes massless irreducible higher spins in AdS space which correspond to the Young tableaux with one row in complete correspondence with [3]. The same result concerning non-uniqueness of the BRST charge will take place if we apply above construction to the system of constraints (5) – (6) describing the propagation of higher massless integer spin fields in the Minkowski space as well.

1 Conclusions

In this paper we have shown that the BRST charge which corresponds to the system of constraints describing higher massless integer spin fields is not unique. Rather there exists a family of charges even for the case of the flat space background. As a result the form and the number of terms which contain the third and higher degrees of the ghost variables may be different. However the final form of the corresponding gauge invariant Lagrangian and of the gauge transformations are such, that the physical spectrum of the theory does not depend on the particular choice of free parameters entering into the BRST charge. That means that in this particular case the cohomologies of the all family BRST charges given by the equation (26) both for the case of the flat and AdS_D backgrounds are the same. Moreover we conjecture that generically the analogous ambiguity if it is observed for other gauge systems should have no impact on the physical spectrum for the different particular choices of the free parameters. The number of free parameters can be different as well. Let us note also that the nilpotence of the BRST charge for the case of AdS_D background requires the modification of the mass – shell operator, which leads the constraint, giving the correct unitary massless representations of AdS_D group.

It seems interesting to apply this procedure to the more complicated representations of the AdS_D group, when the corresponding representations of the flat space time little group is described for two and more rows. It is interesting also to construct the auxiliary representations of the whole nonlinear algebra (18), (6) in terms of Verma module, as well as to study the possibility for the construction of finite dimensional representations for such kind of algebras.

Acknowledgments The work of the authors was supported in part by the INTAS grant, project No 00-00254 and by the RFBR grants. The work of I.L.B and A.P was also supported in part by the joint DFG-RFBR grant. I.L.B is grateful to the INTAS grant, project No 991-590 for partial support. The work of A.P and M.T was supported in part by Votruba-Blokhintsev (Czech Republic-BLTP JINR) grant.

References

- [1] E.S. Fradkin, G.A. Vilkovisky. Phys.Lett. **B 55**, 224 (1975);
I.A. Batalin, G.A. Vilkovisky. Phys.Lett. **B 69**, 309 (1977).
- [2] A. Pashnev, M. Tsulaia. Mod.Phys.Lett. **A13**, 1853 (1998).
- [3] I.L. Buchbinder, A. Pashnev, M. Tsulaia. Phys.Lett. **B523**, 338 (2001).
- [4] Č. Burdík, A. Pashnev, M. Tsulaia. Mod.Phys.Lett. **A15**, 281 (2000).

- [5] Č. Burdík, A. Pashnev, M. Tsulaia. Mod.Phys.Lett. **A16**, 731 (2001);
Nucl.Phys.Proc.Suppl. **102** (2001) 285, hep-th/0103143.
- [6] C. Fronsdal. Phys.Rev. **D18**, 3624 (1978).
- [7] L. Brink, R. Metsaev, M. Vasiliev. Nucl.Phys. **B586**, 183 (2000).
- [8] J. de Boer, F. Harmsze, T. Tjin. Phys.Rept. **272** 139 (1996).